

Contribution of the Defective *BRCA1*, *BRCA2* and *CHEK2* Genes to the Familial Aggregation of Breast Cancer: a Simulation Study Based on the Swedish Family-Cancer Database

Justo Lorenzo Bermejo¹, Alfonso García Pérez², Kari Hemminki^{1,3}

¹Division of Molecular Genetic Epidemiology, German Cancer Research Centre (DKFZ), Heidelberg, Germany; ²Department of Statistics, Spanish Open University (UNED), Madrid, Spain; ³Department of Biosciences at Novum, Karolinska Institute, Huddinge, Sweden

Key words: Familial relative risk, *BRCA1*, *BRCA2*, *CHEK2*, disease penetrance, demographic factors

Corresponding author: Justo Lorenzo Bermejo, Division of Molecular Genetic Epidemiology, German Cancer Research Centre (DKFZ), Im Neuenheimer Feld 580, D-69120, Heidelberg, Germany, e-mail: J.Lorenzo@dkfz.de

Submitted:

Accepted:

Abstract

The known breast cancer susceptibility genes only account for 20% to 25% of the excess familial risk of the disease [1]. The present study assessed the contribution of *BRCA1/2* mutations and *CHEK2* variants to the relative risk of breast cancer for women with affected mothers or sisters. The familial relative risks were estimated by Poisson regression based on the Swedish Family-Cancer Database. The Database was also used to calculate the distribution of life expectancy, the number of daughters per family and the age specific cumulative risk of female breast cancer. This information, together with the penetrances of *BRCA1*, *BRCA2* and *CHEK2* from the literature, was used to simulate the familial clustering of breast cancer under different scenarios. The excess risk explained by *BRCA1*, *BRCA2* and *CHEK2* decreased steeply with the age at diagnosis of the cancers. Around 40% of the familial risk for cases diagnosed before the age of 50 years was associated with *BRCA1/2* mutations. In contrast, roughly 85% of the familial risk of breast cancer diagnosed before the age of 69 years remained unexplained. The contribution of *CHEK2* to familial breast cancer was small.

Introduction

Breast cancer aggregates in families, the disease being about twice as common in mothers and sisters of cases as it is in the general population [2]. The higher risks of breast cancer for monozygotic than for dizygotic twins of cases suggest that the familial aggregation of breast cancer is mainly due to genetic effects, rather than to shared environmental factors [3]. Germline mutations in *BRCA1* and *BRCA2* are frequently found in families containing multiple individuals affected by breast cancer [4]. However,

BRCA1 and *BRCA2* mutations are only identified in about 15-20% of multiple-case families affected by breast cancer alone [5]. *CHEK2**1100delC, a truncating variant that abrogates the kinase activity of *CHEK2* [6], has been also found to contribute significantly to the familial clustering of breast cancer [7]. The variant has shown a frequency of 1.1% in healthy individuals and it has been associated with a breast cancer risk ratio of 1.7 in families without *BRCA1/2* mutations. By contrast, the variant conferred no increased cancer risk in carriers of *BRCA1/2* mutations. The low proportion of familial breast

cancers attributable to known genes, from 20% to 25% [1], reflects major gaps in our knowledge of the genetic background of familial breast cancer.

In addition to the age, sex and genotype specific penetrance, the family history of breast cancer is influenced by demographic factors such as family size and mortality [8]. The aim of the present study was to assess the contribution of the *BRCA1/2* mutations and *CHEK2* variants to the relative risk of breast cancer for women with affected mothers or sisters. We used the Swedish Family-Cancer (SFC) Database to estimate the distribution of life expectancy, the number of daughters per family and the age specific cumulative risk of female breast cancer in Sweden. The penetrances associated with *BRCA1/2* and *CHEK2* were taken from the literature. This information was used to simulate the familial clustering of breast cancer under different scenarios.

Patients and methods

The Swedish Family-Cancer (SFC) Database was created in mid 1990s by linking census information, death notifications and the administrative family registry at Statistics Sweden to the Swedish Cancer Registry. The Database was updated at the end of 2002 to include more than 10.34 million individuals born in Sweden after 1931 as well as more than 810,000 invasive cancers diagnosed after 1958. The Swedish Cancer Registry is based on separate compulsory notifications of cases from clinicians/pathologists or cytologists and is considered to have completeness close to 100% [9]. The incidence of cancer in the Database is similar to the incidence in the Cancer Registry [10, 11]. Data on parity were complete, information on the socioeconomic index and the region was based on population censuses from 1960, 1970, 1980 and 1990. The age of the women in the first generation (mothers) was unrestricted, but the maximum age of women in the second generation (daughters) was 68 years. The present study included 20,742 cases of invasive breast cancer among 3.25 million daughters and 67,575 cases of invasive breast cancer among 2.23 million mothers.

Poisson regression

The relative risk of breast cancer for daughters (RR_{mother}) and sisters (RR_{sister}) of Swedish women affected by breast cancer was estimated by Poisson regression. The women in the SFC Database were followed from birth, immigration date or 1961, whichever came latest, until diagnosis of breast cancer, death,

emigration date or 31 December 2000. The incidence of breast cancer was explained by the variables age (5 year bands), period (10 year bands), parity (six groups from 'any parturition' to 'more than five parturitions'), socioeconomic status (six groups), age at first birth (5 groups, five years bands between 'before age of 20' and 'after age of 35') and residential area (four groups). The analyses were carried out for different restrictions of the age at diagnosis of breast cancer, which varied from 50 to 69 years. Computations were performed with the SAS software using the procedure GENMOD.

Simulation

The SFC Database was used to estimate the cumulative risks of female breast cancer before specific ages. The incidences in *BRCA1/2* mutation carriers reported by Antoniou et al [12] were transformed into cumulative risks by the formula: cumulative risk (%) = $100 \cdot (1 - \exp[-0.05 \cdot \sum x_i])$, where $\sum x_i$ was the sum of the five-year incidences before the age under consideration. The cumulative risks before the age of 69 years were calculated by linear interpolation. The cumulative risks from the SFC Database, the cumulative risks for *BRCA1/2* mutation carriers from the literature and the prevalences of *BRCA1/2* mutations found by Loman et al among affected Swedish women [13] were used to estimate the frequency of *BRCA1/2* mutations in Sweden. For example, the cumulative risk of female breast cancer before the age of 40 years of 0.31%, the cumulative risk of breast cancer by the age of 40 years in *BRCA1* mutation carriers of 11.57% and the prevalence of *BRCA1* mutation carriers in women affected by breast cancer before the age of 40 years of 7.26%, would result in a frequency of *BRCA1* mutations in Sweden of 0.098%. The prevalence and penetrance of *CHEK2* variants were based on the study of the *CHEK2* Breast Cancer Consortium [7]. The distribution of the number of daughters per family and the distribution of life expectancy were calculated using the SFC Database.

The distribution of family size was used to generate one hundred million nuclear families. The genotypes of the parents were created by using the calculated prevalences of *BRCA1/2* mutations and the frequency of *CHEK2* variants from the literature. One allele was taken at random from each parent in order to simulate the genotypes of the daughters, under the assumption that women carrying two copies of one mutated genes were nonviable. The individual's age at death was generated by using the distribution of life expectancy

from the SFC Database. The phenotype of each woman (affected or unaffected) was conditional on her genotype and her age at death. The familial aggregation of breast cancer was explored under different scenarios. The simplest scenario included only one gene; the most elaborated model considered simultaneously BRCA1, BRCA2 and CHEK2. The simulated disease phenotypes were used to calculate the relative risks for daughters and sisters of affected women.

The proportion of familial relative risk attributable to BRCA1, BRCA2 and CHEK2 was assessed by comparing the results from the Poisson regression with the data from the simulation. The formula: $100 \cdot (RR_{BRCA1/2,CHEK2} - 1) / (RR_{mother} - 1)$ was used to calculate the percentage of maternal excess risk attributable to the three genes, where $RR_{BRCA1/2,CHEK2}$ was the estimated relative risk for daughters of affected women when the simulation included the BRCA1, BRCA2 and CHEK2 genes, and RR_{mother} was the relative risk for daughters of affected women estimated by the Poisson regression based on the SFC Database. Similar calculations were carried out to assess the contribution of BRCA1, BRCA2 and CHEK2 to the relative risk for sisters of affected women.

Results

The cumulative risks of breast cancer used in the simulation are presented in Table 1. Based on the SFC

Database, 0.003% of the women had breast cancer by the age of 25 years and 6.48% of them were affected before the age of 69 years. The penetrances of BRCA1/2 mutations estimated by Antoniou et al [12] are also shown in Table 1. These were 63% for women who carried BRCA1 mutations and 42% for BRCA2 mutations carriers by the age of 69 years. The distribution of the number of daughters in the SFC Database was as follows: 68% of the families had one daughter, 26% had two daughters, 5% had three daughters and 1% of the families had four or more daughters. The calculated distribution of life expectancy is shown in Table 1; 77.3% of the women reached the age of 69 years. The cumulative risks of breast cancer from the SFC Database and previous data from the literature resulted in an estimated prevalence of BRCA1 mutations in Sweden of 0.098%, whereas the estimated prevalence of BRCA2 mutations was 0.052%. Following the study of the CHEK2 Breast Cancer Consortium, 1.1% of the simulated individuals were CHEK2 mutation carriers [7]. These data were taken into account to generate the phenotype distribution of a large population, which was utilized to compute the familial risk of breast cancer under different scenarios.

The relative risks of breast cancer for women with affected mothers based on the SFC Database are shown in Fig. 1. The RR_{mother} was 2.11 (95%CI: 1.85-2.41) for breast cancers diagnosed before the age of 50 years, and it decreased to 1.56 (95%CI: 1.46-1.66)

Table 1. Cumulative risk of breast cancer in Sweden, penetrance of BRCA1 and BRCA2 mutations based on the literature and life expectancy of Swedish women*

by age	all women	Cumulative risk of breast cancer (%)		% alive
		BRCA1 mutation carriers	BRCA2 mutation carriers	
25	0.003	0.10	0.10	99.1
30	0.02	0.65	0.70	98.8
35	0.09	4.26	2.47	98.4
40	0.31	11.57	6.20	97.7
45	0.85	23.59	10.37	96.5
50	1.74	38.31	16.18	94.3
55	2.95	45.96	23.05	90.5
60	4.26	53.51	30.55	85.9
65	5.59	59.38	38.61	80.5
69	6.48	62.73	42.25	77.3

*Ref 12; see Patients and methods for details

for cancers under the age of 65 years. The scenario 'cancer occurs due to *CHEK2* mutations' led to estimates of the RR_{mother} practically identical to unity. When only *BRCA2* mutations were considered, the estimated RR_{mother} decreased from 1.04 (breast cancers under the age of 50 years) to 1.01 (cancers under the age of 69 years). The scenario 'cancer is attributable *BRCA1* mutations' showed the RR_{mother} of 1.43 (cancers diagnosed before the age of 50 years) and the RR_{mother} of 1.07 (cancers before the age of 69 years). The maternal risks under the scenario '*BRCA1* and *BRCA2* mutations' were similar to those after the simultaneous consideration of *BRCA1*, *BRCA2* and *CHEK2* mutations; the RR_{mother} was 1.47 for breast cancer before the age of 50 years and 1.09 for breast cancer before the age of 69 years.

The RR_{sister} from the Poisson regression decreased from 2.19 (95%CI: 1.98-2.43), for breast cancers diagnosed before the age of 50 years, to 1.97 (95%CI: 1.86-2.09), for cancers diagnosed before the age of 60 years (Fig. 2). The scenario 'only *BRCA2* mutations' resulted in the RR_{sister} of 1.02 (breast cancers diagnosed before the age of 50 years) and the RR_{sister} of 1.01 (cancers under the age of 69 years). *BRCA1*

mutations showed the RR_{sister} of 1.41 for breast cancers diagnosed before the age of 50 years, and the RR_{sister} of 1.06 for cancers before the age of 69 years. The results for *BRCA1* and *BRCA2* mutations were similar to the results for *BRCA1*, *BRCA2* and *CHEK2*; the RR_{sister} was 1.46 for breast cancer diagnosed before the age of 50 years and it was 1.08 for cancers before the age of 69 years.

The excess risk explained by *BRCA1*, *BRCA2* and *CHEK2* decreased with increasing ages at diagnosis and it was higher for daughters than for sisters of affected women. For example, 14% of the relative risk for daughters of women affected by the age of 69 years was related to *BRCA1*, *BRCA2* and *CHEK2* mutations, but the corresponding proportion for sisters of affected women was only around 8%.

Discussion

The most direct way to address the question concerning the existence of other breast cancer susceptibility genes is to ask whether the known genes can explain the observed familial aggregation of breast cancer [1]. Germline mutations in the *p53*,




Fig. 1. Relative risk of breast cancer for daughters of women with breast cancer in Sweden and effect of *BRCA1*, *BRCA2* and *CHEK2* mutations on the relative risk of breast cancer for daughters of affected women. Both the cases and the probands are restricted to the indicated age

PTEN, *STK11/LKB1* and *ATM* genes are rare in familial breast cancer [14]. The present study assessed the contribution of *BRCA1*, *BRCA2* and *CHEK2* to the relative risk of breast cancer. The estimated familial risks for Swedish women relied entirely on registered data of complete coverage. Other important features of this study were the large number of cases analyzed, the standardization for parity and age of first birth, and the inclusion of information on family size and life expectancy in the simulation. The age of the individuals from the first generation in the SFC Database was unrestricted, but the maximum age of the individuals in the second generation (68 years) was a limitation on the present study.

Mutations in *BRCA1* and *BRCA2* show considerable ethnic and geographic variation [15]. Specific *BRCA1* or *BRCA2* mutations have become common as a result of founder effects in Ashkenazi Jewish populations [15-17], Poland [18], Iceland [19] and the European part of Russia [20]. The contribution of *BRCA1* and *BRCA2* to familial breast cancer in those populations is likely to be more important than in Sweden. The mutation prevalences assumed in this study, 0.098% for *BRCA1* and 0.052% for *BRCA2*, were calculated using the SFC Database and

the Swedish results of Lohman et al [13]. These prevalences are in agreement with the literature, e.g. Easton proposed frequencies between 0.05% and 0.20% for both *BRCA1* and *BRCA2* [1] and Domchek et al estimated a prevalence of *BRCA1* mutations of 0.125% [21]. The prevalence and penetrance assumed for *CHEK2* variants may be more inaccurate. This study concentrated on the 1100delC frameshift mutation and was based on the results from the *CHEK2*-Breast Cancer Consortium, but the same conclusions were reached by using German [22] or Finish [23] data (results not shown).

The estimated risks were higher for sisters than for daughters of affected women, and the difference between the two familial relative risks increased with the age at diagnosis of breast cancer. Several studies have suggested that this difference is mostly attributable to the unequal number of parturitions and the different calendar year of diagnosis of mothers and sisters [2, 24]. Since 97% of the analyzed daughters were parous, the contribution of parity to the difference of relative risks, if any, should be small. In contrast, the median calendar year of diagnosis of mothers was 1971 and that of sisters - 1996. The establishment of screening services in Sweden has resulted in the earlier detection




Fig. 2. Relative risk of breast cancer for sisters of women with breast cancer in Sweden and effect of *BRCA1*, *BRCA2* and *CHEK2* mutations on the relative risk of breast cancer for sisters of affected women. Both the cases and the probands are restricted to the indicated age

of breast cancers [25] and may have also affected the detection of familial cancers.

In contrast to the familial relative risks estimated by the Poisson regression, the relative risks associated with *BRCA1* and *BRCA2* mutations in the simulation were slightly lower for sisters than for mothers of affected women, especially when the cancers were diagnosed before the age of 50 years. The separate analysis of each gene showed that *BRCA1* mutations would explain 34% to 39% of the familial relative risk by the age of 50 years, the corresponding figure would be 2% to 4% for *BRCA2* mutations, and less than 1% of the familial risk was attributable to *CHEK2* variants.

In conclusion, the proportion of excess familial risk due to *BRCA1/2* mutations cited in the literature, about 15% [26], varies considerably with the population and the age at diagnosis of the cancers. In Sweden roughly 40% of the familial relative risk for breast cancers diagnosed before the age of 50 years is likely to be associated with *BRCA1/2* mutations, but around 85% of the excess risk remains unexplained when all cancers diagnosed before the age of 69 years are considered. The proportion of familial excess attributable to *CHEK2* variants, or other low susceptibility genes, is small.

Acknowledgements

The study was supported by Deutsche Krebshilfe and the Swedish Cancer Society. The Family-Cancer Database was created by linking registers maintained by Statistics Sweden and the Swedish Cancer Registry.

References

1. Easton DF. How many more breast cancer predisposition genes are there? *Breast Cancer Res* 1999; 1: 14-17.
2. Pharoah PD, Day NE, Duffy S, Easton DF and Ponder BA. Family history and the risk of breast cancer: a systematic review and meta-analysis. *Int J Cancer* 1997; 71: 800-809.
3. Lichtenstein P, Holm NV, Verkasalo PK, Iliadou A, Kaprio J, Koskenvuo M, Pukkala E, Skytthe A and Hemminki K. Environmental and heritable factors in the causation of cancer—analyses of cohorts of twins from Sweden, Denmark, and Finland. *N Engl J Med* 2000; 343: 78-85.
4. Ford D, Easton DF, Stratton M, Narod S, Goldgar D, Devilee P, Bishop DT, Weber B, Lenoir G, Chang-Claude J, Sobol H, Teare MD, Struwing J, Arason A, Scherneck S, Peto J, Rebbeck TR, Tonin P, Neuhausen S, Barkardottir R, Eyfjord J, Lynch H, Ponder BA, Gayther SA, Zelada-Hedman M, et al. Genetic heterogeneity and penetrance analysis of the *BRCA1* and *BRCA2* genes in breast cancer families. The Breast Cancer Linkage Consortium. *Am J Hum Genet* 1998; 62: 676-689.
5. Antoniou AC, Pharoah PD, McMullan G, Day NE, Ponder BA and Easton D. Evidence for further breast cancer susceptibility genes in addition to *BRCA1* and *BRCA2* in a population-based study. *Genet Epidemiol* 2001; 21: 1-18.
6. Wu X, Webster SR and Chen J. Characterization of tumor-associated *Chk2* mutations. *J Biol Chem* 2001; 276: 2971-2974.
7. Meijers-Heijboer H, van den Ouweland A, Klijn J, Wasielewski M, de Snoo A, Oldenburg R, Hollestelle A, Houben M, Crepin E, van Veghel-Plandsoen M, Elstrodt F, van Duijn C, Bartels C, Meijers C, Schutte M, McGuffog L, Thompson D, Easton D, Sodha N, Seal S, Barfoot R, Mangion J, Chang-Claude J, Eccles D, Eeles R, Evans DG, Houlston R, Murday V, Narod S, Peretz T, Peto J, Phelan C, Zhang HX, Szabo C, Devilee P, Goldgar D, Futreal PA, Nathanson KL, Weber B, Rahman N and Stratton MR. Low-penetrance susceptibility to breast cancer due to *CHEK2*(*)1100delC in noncarriers of *BRCA1* or *BRCA2* mutations. *Nat Genet* 2002; 31: 55-59.
8. Sibert A and Goldgar DE. The effect of disease penetrance, family size, and age of onset on family history with application to setting eligibility criteria for genetic testing. *Fam Cancer* 2003; 2: 35-42.
9. Cancer incidence in Sweden, 2000. Stockholm, National Board of Health and Welfare 2002; pp. 118.
10. Hemminki K, Li X, Plna K, Granstrom C and Vaitinen P. The nation-wide Swedish family-cancer database—updated structure and familial rates. *Acta Oncol* 2001; 40: 772-777.
11. Plna K and Hemminki K. Familial bladder cancer in the National Swedish Family Cancer Database. *J Urol* 2001; 166: 2129-2133.
12. Antoniou A, Pharoah PD, Narod S, Risch HA, Eyfjord JE, Hopper JL, Loman N, Olsson H, Johannsson O, Borg A, Pasini B, Radice P, Manoukian S, Eccles DM, Tang N, Olah E, Anton-Culver H, Warner E, Lubinski J, Gronwald J, Gorski B, Tulinius H, Thorlacius S, Eerola H, Nevanlinna H, Syrjakoski K, Kallioniemi OP, Thompson D, Evans C, Peto J, Lalloo F, Evans DG and Easton DF. Average risks of breast and ovarian cancer associated with *BRCA1* or *BRCA2* mutations detected in case Series unselected for family history: a combined analysis of 22 studies. *Am J Hum Genet* 2003; 72: 1117-1130.
13. Loman N, Bladstrom A, Johannsson O, Borg A and Olsson H. Cancer incidence in relatives of a population-based set of cases of early-onset breast cancer with a known *BRCA1* and *BRCA2* mutation status. *Breast Cancer Res* 2003; 5: R175-186.
14. Arver B, Du Q, Chen J, Luo L and Lindblom A. Hereditary breast cancer: a review. *Semin Cancer Biol* 2000; 10: 271-288.
15. Neuhausen SL. Ethnic differences in cancer risk resulting from genetic variation. *Cancer* 1999; 86: 2575-2582.
16. Struwing JP, Hartge P, Wacholder S, Baker SM, Berlin M, McAdams M, Timmerman MM, Brody LC and Tucker MA. The risk of cancer associated with specific mutations of *BRCA1* and *BRCA2* among Ashkenazi Jews. *N Engl J Med* 1997; 336: 1401-1408.
17. Tonin P, Ghadirian P, Phelan C, Lenoir GM, Lynch HT, Letendre F, Belanger D, Monte M and Narod SA. A large multisite cancer family is linked to *BRCA2*. *J Med Genet* 1995; 32: 982-984.
18. Gorski B, Jakubowska A, Huzarski T, Byrski T, Gronwald J, Grzybowska E, Mackiewicz A, Stawicka M, Bebenek M, Sorokin D, Fiszera-Maliszewska L, Haus O, Janiszewska H, Niepsuj S, Gozdz S, Zaremba L, Posmyk M, Pluzanska M, Kilar E, Czudowska D, Wasko B, Miturski R, Kowalczyk JR, Urbanski K, Swiec M, Koc J, Debniak B, Rozmiarek A, Debniak T, Cybulski C, Kowalska E, Toloczko-Grabarek A, Zajaczek S, Menkiszak J, Medrek K, Masojc B, Mierzejewski M, Narod SA and Lubinski J. A high proportion of founder *BRCA1* mutations in Polish breast cancer families. *Int J Cancer* 2004; 110: 683-686.
19. Thorlacius S, Olafsdottir G, Tryggvadottir L, Neuhausen S, Jonasson JG, Tavtigian SV, Tulinius H, Ogmundsdottir HM and

- Eyford JE. A single BRCA2 mutation in male and female breast cancer families from Iceland with varied cancer phenotypes. *Nat Genet* 1996; 13: 117-119.
20. Loginova AN, Pospekhova NI, Lyubchenko LN, Budilov AV, Zakhar'ev VM, Gar'kavtseva RF, Ginter EK and Karpukhin AV. Spectrum of mutations in BRCA1 gene in hereditary forms of breast and ovarian cancer in Russian families. *Bull Exp Biol Med* 2003; 136: 276-278.
 21. Domchek SM, Eisen A, Calzone K, Stopfer J, Blackwood A and Weber BL. Application of breast cancer risk prediction models in clinical practice. *J Clin Oncol* 2003; 21: 593-601.
 22. Dufault MR, Betz B, Wappenschmidt B, Hofmann W, Bandick K, Golla A, Pietschmann A, Nestle-Kramling C, Rhiem K, Huttner C, von Lindern C, Dall P, Kiechle M, Untch M, Jonat W, Meindl A, Scherneck S, Niederacher D, Schmutzler RK and Arnold N. Limited relevance of the CHEK2 gene in hereditary breast cancer. *Int J Cancer* 2004; 110: 320-325.
 23. Vahteristo P, Bartkova J, Eerola H, Syrjakoski K, Ojala S, Kilpivaara O, Tamminen A, Kononen J, Aittomaki K, Heikkila P, Holli K, Blomqvist C, Bartek J, Kallioniemi OP and Nevanlinna H. A CHEK2 genetic variant contributing to a substantial fraction of familial breast cancer. *Am J Hum Genet* 2002; 71: 432-438.
 24. Easton DF, Matthews FE, Ford D, Swerdlow AJ and Peto J. Cancer mortality in relatives of women with ovarian cancer: the OPCS Study. Office of Population Censuses and Surveys. *Int J Cancer* 1996; 65: 284-294.
 25. Zahl PH, Strand BH and Maehlen J. Incidence of breast cancer in Norway and Sweden during introduction of nationwide screening: prospective cohort study. *BMJ* 2004; 328: 921-924.
 26. Easton DF. Familial risks of breast cancer. *Breast Cancer Res* 2002; 4: 179-181.